

Product datasheet

RAP, rat recombinant, 250 µg

Short overview

Cat. No.	62021
Quantity	250 µg

Product description

Source	Rat, recombinant (his:RAP:c-myc fusion protein)
Molecular Weight	40 kDa
Purity	> 95% (determined by SDS gelelectrophoresis)
Reconstitution	When reconstituted with 250 µl distilled water, solution will contain 1 mg/ml rat RAP protein in TBS, pH 7.5, 0.1% BSA, 0.09% NaN ₃ .
Application	Protein standard in 1D and 2D SDS gelelectrophoresis, immunoblotting and receptor-binding studies
Storage	Lyophilized at 2-8°C; reconstituted at -20°C (avoid freeze/thaw cycles)
Intended use	Research use only

Background

Protein standard for immunoblotting, immunization, immunoassay and receptor-binding studies. Sequence highly homolog with human RAP. Rat his:RAP:c-myc fusion protein. Ligand binding to RAP is Ca²⁺ dependent and e.g. lipid receptors can be released from RAP by a buffer containing 10 mM EDTA (Bajari et al. 2005). Furthermore, buffers containing phosphate should be avoided (it would form precipitates with Ca²⁺).

Product images



RAP, rat recombinant, 250 µg

References

Publication	Species	Application
Hayashi, H. Glial Lipoproteins Stimulate Axon Growth of Central Nervous System Neurons in Compartmented Cultures. 14 (2004).		
Bajari, T. M. et al. A Minimal Binding Domain of the Low Density Lipoprotein Receptor Family. 379, 1053-1062 (1998).		