

Product datasheet

anti-Lewis a mouse monoclonal, 7LE, purified

Short overview

Cat. No.	691517
Quantity	1 ml (100 µg/ml)
Concentration	100 µg/ml

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1 kappa
Clone	7LE
Immunogen	Ovarian cyst mucin
Formulation	PBS with 0.02% sodium azide
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	2-8°C
Intended use	Research use only
Application	ELISA, FACS, ICC/IF, IHC, WB
Reactivity	Human, Mouse

Applications


ELISA	Assay dependent
Flow Cytometry (FACS)	0.5-1.0 µg/million cells in 0.1 ml
Immunocytochemistry (ICC)	1:100-1:200 (0.5-1.0 µg/ml)
Immunohistochemistry (IHC) - frozen	1:50-1:100 (1-2 µg/ml)
Immunohistochemistry (IHC) - paraffin	1:50-1:100 (1-2 µg/ml; microwave treatment in 10 mM citrate buffer pH 6.0 recommended)
Western Blot (WB)	1:50-1:100 (1-2 µg/ml)

Background

7LE reacts with Lewis a blood group antigen, a carbohydrate determinant present on both glycolipids and glycoproteins, expressed a.o. in colonic and kidney epithelial cells. Lewis a may be useful for detection of gastrointestinal, pancreatic and colorectal carcinomas.

Positive control: human colon.

Product images



Human colon