

## **Product datasheet**

# anti-Desmocollin 1 mouse monoclonal, Dsc1-U100, liquid, purified, sample

#### Short overview

 Cat. No.
 690092S

 Quantity
 200 µl

Concentration 50 μg/ml (10 μg)

#### **Product description**

HostMouseAntibody TypeMonoclonalIsotypeIgG1CloneDsc1-U100

Immunogen Synthetic peptide corresponding to sequence present within the intracellular part of human

desmocollin 1 (HQTLESVKGVGQGDTGRYATDWQC)

**Formulation** PBS pH 7.4 with 0.09% sodium azide and 0.5% BSA

UniprotID Q08554 (Human)

**Synomym** Desmocollin-1, DSC1, Cadherin family member 1, Desmosomal glycoprotein 2/3, DG2/DG3,

CDHF1

Conjugate Unconjugated

**Purification** Affinity chromatography

Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles

Intended use Research use only

Application IHC, WB

Reactivity Human, Mouse, Rat

#### **Applications**

Immunohistochemistry (IHC) - frozen 1:50-1:100 (0.5-1 µg/ml, preincubation (directly after fixation) with

0.05-0.2% Triton X-100, for 5-10 min, depending on tissue type, is recommended; without detergent pretreatment a cytoplasmic

component might cross-react)

**Immunohistochemistry (IHC) - paraffin** 1:50-1:100 (0.5-1 μg/ml, microwave treatment recommended; a slight

crossreactivity with a lymphocyte subpopulation is abolished by mild

trypsinization after the microwave step)

Western Blot (WB) Assay dependent

#### Background


Mouse monoclonal antibody localizes Dsc1 present in

- Suprabasal layers of interfollicular epidermis
- Specific cell layers (e.g. in hair follicle root sheath)
- Hassall bodies (spheroidal formations of densely packed reticulum-derived cells in the thymus).


Reacts with 111 kDa (form "a") and 103 kDa (form "b") polypeptides (desmocollin 1 splice forms, both at pl 5.2) from human epidermal desmosomes.

Reactivity on cultured cell lines: positive on primary keratinocytes, however, restricted to local piles of differentiated cells. All simple epithelia are negative.


### **Product images**



IHC of rat skin with anti-desmocollin 1 (1:50, courtesy of J.Heß, University Hospital Heidelberg)



anti-Desmocollin 1 mouse monoclonal, Dsc1-U100, liquid, purified, sample



IHC of mouse skin with anti-desmocollin 1 (1:50, courtesy of J.Heß, University Hospital Heidelberg)

## References

Publication	Species	Application
Akat, K. et al. Molecular characterization of desmosomes in	human	IHC (frozen)
meningiomas and arachnoidal tissue. Acta Neuropathol. 106,		
<u>337–347 (2003)</u> .		
Kurzen, H. et al. Compositionally different desmosomes in the	human	IHC (paraffin)
various compartments of the human hair follicle. Differentiation		
<u>63, 295–304 (1998).</u>		