

Product datasheet

anti-Cardiac Actin mouse monoclonal, AC1-20.4.2, liquid, purified, sample

Short overview

Cat. No.	690075S
Quantity	200 µl
Concentration	50 µg/ml (10 µg)

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1
Clone	AC1-20.4.2
Immunogen	Synthetic NH2 terminus decapeptide of cardiac isoform of actin
Formulation	PBS pH 7.4 with 0.09% sodium azide and 0.5% BSA
UniprotID	P68034 (Chicken), P68032 (Human), G1STB6 (Rabbit)
Synonym	Actin, alpha cardiac muscle 1, Alpha-cardiac actin [Cleaved into: Actin, alpha cardiac muscle 1, intermediate form], ACTC1, ACTC
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
Intended use	Research use only
Application	IHC, WB
Reactivity	Bovine, Chicken, Human, Mouse, Rabbit


Applications

Immunohistochemistry (IHC) - frozen	1:10 (5 µg/ml; include 0.5 M NaCl in all washing buffers to enhance specificity)
Immunohistochemistry (IHC) - paraffin	1:10 (5 µg/ml; microwave treatment recommended)
Western Blot (WB)	1:1,000 (50 ng/ml; include 1 M NaCl in all washing buffers to enhance specificity)


Background

Ac1 represents an excellent marker for cardiac tissue; it discriminates fetal (cardiac) a-actin from all other actin isoforms. Fetal actin can be localized in regenerating skeletal muscle after injury (in satellite cells) and in veins of the umbelical cord. Mab Ac1-20.4.2 shows no cross reaction with other actin isoforms present in skeletal and smooth muscle, provided that stringent experimental conditions have been applied. Polypeptide reacting: Specific for fetal (cardiac) isoform of actin.

Product images



anti-Cardiac Actin mouse monoclonal, AC1-20.4.2, liquid, purified, sample



Western blot analysis of human heart whole tissue lysate and with anti-Cardiac Actin antibody. Western blot analysis was performed on 20 µg human heart lysate. The PVDF membrane was blocked with 5% milk in PBST (PBS + 0.1% Tween 20) for 1 h at RT. The primary antibody anti-Cardiac Actin mouse monoclonal, AC1-20.4.2 (Cat. No. 690075) was diluted in blocking buffer (antibody concentration 0.05 µg/ml) and incubated for 1 h at RT. The secondary antibody anti-mouse IgG, HRP conjugate was also diluted in blocking buffer (antibody concentration 0.2 µg/ml) and incubated for 1 h at RT. The bands were visualized by chemiluminescent detection using Pierce™ ECL Western Blotting Substrate.

References

Publication	Species	Application
Franke, W. W. et al. Specific immunohistochemical detection of cardiac/fetal α-actin in human cardiomyocytes and regenerating skeletal muscle cells. Differentiation 60, 245-250 (1996).	human	IHC (frozen), IHC (paraffin)
Rudy, D. E., Yatskievych, T. A., Antin, P. B. & Gregorio, C. C. Assembly of thick, thin, and titin filaments in chick precardiac explants. Dev. Dyn. 221, 61-71 (2001).	chicken	ICC-IF
Domke, L and Franke, W. The cell-cell junctions of mammalian testes... Cell Tissue Res, 375, 451-482, (2019)	bovine	ICC-IF
Lindskog, C. et al. The human cardiac and skeletal muscle proteomes defined by transcriptomics and antibody-based profiling. BMC Genomics 16, 475 (2015).	human	IHC
Marrocco, V. et al. PKC and PKN in heart disease. J.Mol.Cell.Cardiol. 128, 212-226 (2019)	mouse	WB