

Product datasheet

anti-Keratin K7 mouse monoclonal, Ks7.18, prediluted, purified

Short overview

Cat. No.	65025
Quantity	5 ml

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1
Clone	Ks7.18
Immunogen	Cytoskeletal proteins from cultured HeLa cells
Formulation	PBS pH 7.4 with 0.5% BSA and 0.09% sodium azide
UniprotID	Q29S21 (Bovine), P08729 (Human), A0A287ASI0 (Pig), H0VIA2 (Guinea pig), A0A6P7DW90 (Sheep)
Synonym	Keratin, type II cytoskeletal 7, Cytokeratin-7, CK-7, Keratin-7, K7, Sarcolectin, Type-II keratin Kb7, KRT7, SCL
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
Intended use	Research use only
Application	ICC/IF, IHC, WB
Reactivity	Bovine, Human, Pig, Sheep
No reactivity	Dog, Mouse, Rabbit, Rat

Applications

Immunocytochemistry (ICC)	Assay dependent
Immunohistochemistry (IHC) - frozen	Ready-to-use
Immunohistochemistry (IHC) - paraffin	Ready-to-use (protease treatment and/ or microwave treatment recommended)
Western Blot (WB)	Assay dependent

Background

Ks7.18 represents an excellent marker for the discrimination of specific subtypes of adenocarcinoma: e.g. adenocarcinoma of pancreas, bile duct carcinoma and transitional carcinoma of bladder are stained, whereas hepatocellular and prostate carcinomas are negative. Detects specific subtypes of adenocarcinomas: adenocarcinoma of pancreas, gallbladder, lung, cervix; cholangio carcinoma of liver; ductal and lobular carcinoma of breast; carcinomas of ovary; transitional cell carcinoma of bladder; mesothelioma; negative with most cases of hepatocellular carcinoma. In colorectal carcinoma early stages are reported to be negative but advanced stages of tumor development are positive for keratin K7 expression. Occasionally, staining of blood vessel walls, particularly of endothelial cells may be observed. Reacts with Mr 54,000 polypeptide (keratin K7;

PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg


Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: info@progen.com | Web: www.progen.com

2024 April 20 / Version: 65025/DS-190221ibg | Page 1


formerly also designated cytokeratin 7) of human glandular epithelia.

Reactivity on cultured cell lines: HeLa, RT 112, T24, BT-20, CAMA-1, Detroit 562, (MCF-7 and HT-29 are negative).


Product images



Human thyroid carcinoma (courtesy of J.Heß, University Hospital Heidelberg)



Human kidney (courtesy of J.Heß, University Hospital Heidelberg)



Human kidney (courtesy of J.Heß, University Hospital Heidelberg)

References

Publication	Species	Application
Demirkesen, C., Hoede, N. & Moll, R. Epithelial markers and differentiation in adnexal neoplasms of the skin: an immunohistochemical study including individual cytokeratins. J. Cutan. Pathol. 22, 518â€“535 (1995).	human	IHC (paraffin)
Langbein, L., Yoshida, H., Praetzel-Wunder, S., Parry, D. A. & Schweizer, J. The Keratins of the Human Beard Hair Medulla: The Riddle in the Middle. J. Invest. Dermatol. 130, 55â€“73 (2010).	human	IHC (frozen)
Alam, C. M. et al. Decreased levels of keratin 8 sensitize mice to streptozotocin-induced diabetes.Â Acta Physiol (Oxf).224,e13085(2018).	mouse	IHC (frozen)/IF