

Product datasheet

anti-DDDDK-tag mouse monoclonal, AP1501, lyophilized, purified

Short overview

Cat. No.	910DDK
Quantity	25 µg
Concentration	0.25 mg/ml after reconstitution with 100 µl PBS

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG2b
Clone	AP1501
Immunogen	Recombinant protein containing the sequence DDDDK
Formulation	Lyophilized; reconstitute in 100 µl sterile PBS, pH 7.4
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage before reconstitution	2-8°C until indicated expiry date
Storage after reconstitution	-20°C (avoid freeze/thaw cycles)
Intended use	Research use only
Application	ICC/IF, IP, WB
Reactivity	DDDDK


Applications

Immunocytochemistry (ICC)	Assay dependent
Immunoprecipitation (IP)	Assay dependent
Western Blot (WB)	1:50,000-1:100,000 (0.005-0.0025 µg/ml)

Background

The monoclonal AP1501 antibody recognizes the amino acid sequence DDDDK (Flag). The DDDDK-tag is commonly added to recombinant proteins and can be used for detection or purification of the tagged protein.

Product images



Western blot analysis of E. coli lysate containing DDDDK-tagged protein with anti-DDDDK-tag antibody. Western blot analysis was performed on 80 ng, 40 ng or 10 ng of E. coli lysate containing DDDDK-tagged protein. Cells were lysed with SDS sample buffer. The PVDF membrane was blocked with 5% dry milk in PBST for 1 h at RT. The primary antibody anti-DDDDK-tag mouse monoclonal, AP1501 (Cat. No. 910DDKL) was diluted in blocking buffer (antibody concentration 0.005 µg/ml) and incubated for 1 h at RT. The secondary antibody goat anti-mouse IgG polyclonal, HRP conjugate was also diluted in blocking buffer (antibody concentration 0.2 µg/ml) and incubated for 1 h at RT. The bands were visualized by chemiluminescent detection using Pierce™ ECL Western Blotting Substrate.