

Product datasheet

anti-AAV2 (intact particle) mouse monoclonal, A20, Biotin Conjugate

Short overview

Cat. No.	61555
Quantity	750 µl

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG3
Clone	A20
Immunogen	AAV2 capsids
Formulation	Lyophilized; reconstitute with 750 µl sterile PBS
Synonym	Adeno-associated virus 2; AAV-2
Conjugate	Biotin
Purification	Affinity chromatography
Storage before reconstitution	2-8°C until indicated expiry date
Storage after reconstitution	2-8°C
Intended use	Research use only
Application	ELISA
Reactivity	AAV2, AAV2 7m8, AAV3
No reactivity	AAV1, AAV11, AAV12, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAVDJ, AAVrh10, AAVrh74

Applications

ELISA	1:20
Neutralization Assay	EC50 ~5 ng/ml (AAV2) and ~3 ng/ml (AAV3) - assay dependent


Background

For characterization of different stages of infection and very useful for the analysis of the AAV2 assembly process. A20 specifically reacts with AAV2 and AAV3, empty and full capsids. Recognizes a conformational epitope of assembled capsids, not present in denatured capsid proteins and native but unassembled capsid proteins. The antibody cannot be used for immunoblotting. Epitope mapping experiments (Wobus et al. 2000) identified four immunoreactive (discontinuous) regions. The major reaction was attributed to sequence aa 369 to aa378 of AAV2 capsids.

Wobus, C. E. et al. Monoclonal antibodies against the adeno-associated virus type 2 (AAV-2) capsid: epitope mapping and identification of capsid domains involved in AAV-2-cell interaction and neutralization of AAV-2 infection. J. Virol. 74, 928193 (2000). Moskalenko, M. et al. Epitope Mapping of Human Anti-Adeno-Associated Virus Type 2 Neutralizing Antibodies: Implications for Gene Therapy and Virus Structure. Journal Virol. 74, 17611766 (2000).

Limited Use Label License: Research Use Only Product is exclusively licensed to PROGEN Biotechnik GmbH. The use of these products for the development, manufacturing and sale of secondary products/derivatives which are based on the purchased products and/or which include the purchased product require a royalty based sub-license agreement.

Product images



© PROGEN

ELISA with A20 Biotin Conjugate (Cat. No. 61555) as detection antibody (1:20)

References

Publication	Species	Application
<u>Wistuba, A., Weger, S., Kern, A., Rgen, J. & Kleinschmidt, A. Intermediates of Adeno-Associated Virus Type 2 Assembly: Identification of Soluble Complexes Containing Rep and Cap Proteins. J. Virol. 69, 5311â€“5319 (1995).</u>	AAV2	IP
<u>Moskalenko, M. et al. Epitope mapping of Human Anti-Adeno-Associated Virus Type 2 Neutralizing Antibodies: Implications for Gene Therapy and Virus Structure. Jounal Virol. 74, 1761â€“1766 (2000).</u>	AAV2	neutralization, epitope mapping
<u>McCraw, D. M., Oâ€™Donnell, J. K., Taylor, K. A., Stagg, S. M. & Chapman, M. S. Structure of adeno-associated virus-2 in complex with neutralizing monoclonal antibody A20. Virology 431, 40â€“49 (2012).</u>	AAV2	cryoEM
<u>McCraw, D. M., Oâ€™Donnell, J. K., Taylor, K. A., Stagg, S. M. & Chapman, M. S. Structure of adeno-associated virus-2 in complex with neutralizing monoclonal antibody A20. Virology 431, 40â€“49 (2012).</u>	AAV2	cryoEM
<u>Huttner, N. A. et al. Genetic modifications of the adeno-associated virus type 2 capsid reduce the affinity and the neutralizing effects of human serum antibodies. Gene Ther. 10, 2139â€“2147 (2003).</u>	AAV2	epitope mapping