

Product datasheet

anti-MAP3K1 mouse monoclonal, EBS-T-008, purified

Short overview

Cat. No.	691675
Quantity	1 ml (100 µg/ml)
Concentration	100 µg/ml

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG2a kappa
Clone	EBS-T-008
Immunogen	Partial recombinant MAP3K1 (aa1211-1310)
Formulation	PBS with 0.02% sodium azide
UniprotID	Q13233 (Human)
Synonym	Mitogen-activated protein kinase kinase kinase 1, EC 2.7.11.25, MAPK/ERK kinase kinase 1, MEK kinase 1, MEKK 1, MAP3K1, MAPKKK1, MEKK, MEKK1
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	2-8°C
Intended use	Research use only
Application	ELISA, FACS, ICC/IF, IHC, WB
Reactivity	Human

Applications

ELISA	Assay dependent
Flow Cytometry (FACS)	0.5-1.0 µg/million cells in 0.1 ml
Immunocytochemistry (ICC)	Assay dependent
Immunohistochemistry (IHC) - frozen	1:100-1:200 (0.5-1.0 µg/ml)
Immunohistochemistry (IHC) - paraffin	1:100-1:200 (0.5-1.0 µg/ml; microwave treatment in 10 mM Tris with 1 mM EDTA pH 9.0 recommended)
Western Blot (WB)	1:100-1:200 (0.5-1.0 µg/ml)

Background

MAPKs are involved in directing cellular responses to a diverse array of potentially harmful stimuli, such as mitogens, osmotic stress, heat shock, proinflammatory cytokines, but also growth factors (mammals). Possibly located exclusively in the cell nucleus, they regulate cell functions including proliferation, gene expression, differentiation, mitosis, cell survival, and apoptosis. In order to become active, they require usually multiple phosphorylation events in their activation loops, including phosphorylation by MAP2 kinases (Ste-7 kinases), which in turn are phosphorylated by the MAP3 kinase family, of which many are located at the cell membrane. Thus through this pathway, stimuli can effectively

PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg


Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: info@progen.com | Web: www.progen.com

2024 April 26 / Version: 691675/DS-230420ibg | Page 1


be conveyed from the cell membrane to the nucleus. Inactivation of MAPKs takes place by several phosphorylases, including dedicated phosphorylases.

Positive control: A431, HeLa or HL-60 cells or liver tissue.


Product images



Human uterine carcinoma



Human cervical cancer



Human thyroid carcinoma