

Product datasheet

anti-CD98 mouse monoclonal, IPO-T10, purified

Short overview

Cat. No. 691610

Quantity1 ml (100 μ g/ml)Concentration100 μ g/ml

Product description

Host Mouse
Antibody Type Monoclonal
Isotype IgM kappa
Clone IPO-T10

Immunogen Stimulated human PBL

Formulation PBS with 0.02% sodium azide

UniprotID P08195 (Human)

Synomym 4F2 cell-surface antigen heavy chain, 4F2hc, 4F2 heavy chain antigen, Lymphocyte activation

antigen 4F2 large subunit, Solute carrier family 3 member 2, CD antigen CD98, SLC3A2, MDU1

Conjugate Unconjugated

Purification Affinity chromatography

Storage 2-8°C

Intended use Research use only

Application FACS, IHC **Reactivity** Human

Applications

Flow Cytometry (FACS) 0.5-1.0 μg/million cells in 0.1 ml

Immunohistochemistry (IHC) - frozen 1:50-1:100 (1-2 μg/ml)


Background

CD98 exists as a heterodimer containing a disulphide-linked glycosylated heavy chain and a non-glycosylated light chain. It is a member of the solute carrier family and encodes a cell surface, transmembrane protein. The protein exists as the heavy chain of a heterodimer, covalently bound through disulphide bonds to one of several possible light chains. The encoded transporter plays a role in regulation of intercellular calcium levels and transport L-type amino acids. Alternatively spliced transcript variants, encoding different isoforms, have been characterized.

Monocytes express high levels of CD98 antigen. Peripheral blood T- and B-cells, as well as NK-cells and granulocytes express low levels of CD98. Activation of T-cells and NK-cells leads to upregulation of CD98. RBCs are negative. IPO-T10 was typed at the VIth International Workshop and Conference on Human Leukocyte Differentiation Antigens.

 $Positive\ control:\ Jurkat,\ MG63,\ HUT-78,\ K562,\ YT,\ U937\ and\ human\ lymphocytes\ or\ tonsils.$

Product images



FACS with K562 cells