

## Product datasheet

anti-AAV1 (intact particle) mouse monoclonal, ADK1a, lyophilized, purified

### Short overview

<b>Cat. No.</b>	610150
<b>Quantity</b>	50 µg
<b>Concentration</b>	50 µg/ml after reconstitution with 1 ml dist. water

### Product description

<b>Host</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Isotype</b>	IgG2a lambda
<b>Clone</b>	ADK1a
<b>Immunogen</b>	AAV1 capsids
<b>Formulation</b>	Lyophilized; reconstitute in 1 ml dist. water (final solution contains 0.09% sodium azide, 0.5% BSA in PBS buffer, pH 7.4)
<b>Binding affinity</b>	KD value (AAV1) = <1.0E-12 M KD value (AAV6) = <1.0E-12 M
<b>Synonym</b>	Adeno-associated virus 1; AAV-1
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity chromatography
<b>Storage before reconstitution</b>	2-8°C until indicated expiry date
<b>Storage after reconstitution</b>	Up to 3 months at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
<b>Intended use</b>	Research use only
<b>Application</b>	Affinity chromatography, Dot blot, ELISA, ICC/IF, IP, Neutralization assay
<b>Reactivity</b>	AAV1, AAV12, AAV6
<b>No reactivity</b>	AAV11, AAV2, AAV3, AAV4, AAV5, AAV7, AAV8, AAV9, AAVDJ, AAVrh10, AAVrh74

### Applications

<b>Affinity Chromatography</b>	Assay dependent
<b>Dot Blot</b>	1:500 (0.1 µg/ml; non-denaturing conditions)
<b>ELISA</b>	Assay dependent
<b>Immunocytochemistry (ICC)</b>	1:20
<b>Immunoprecipitation (IP)</b>	1:5
<b>Neutralization Assay</b>	EC50 ~2 ng/ml (AAV1) and ~2 ng/ml (AAV6) - assay dependent

### Background

PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg


Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: [info@progen.com](mailto:info@progen.com) | Web: [www.progen.com](http://www.progen.com)

2024 April 23 / Version: 610150/DS-020124lim | Page 1

For characterization of different stages of infection and very useful for the analysis of the AAV assembly process. ADK1a specifically reacts with intact adeno-associated virus particles, empty and full capsids. Recognizes a conformational epitope of assembled capsids, not present in denatured capsid proteins and native but unassembled capsid proteins. The antibody cannot be used for immunoblotting.

Limited Use Label License: Research Use OnlyProduct is exclusively licensed to PROGEN Biotechnik GmbH. The use of these products for the development, manufacturing and sale of secondary products/derivatives which are based on the purchased products and/or which include the purchased product require a royalty based sub-license agreement.

Product images



Neutralization of AAV1-GFP vectors with the ADK1a antibody (Cat. No. 610150). AAV infection was shown in HeLa cells and photos (GFP, CPE, merge) were taken ~48 h post infection. Neutralization was enhanced with increasing ADK1a concentration.


Serotype	Clone	Contact residues	Footprint residues
AAV1	ADK1a	448 R, 450 Q, <u>453-SGSAQ-457</u> , 500 N	262- <u>#263-265-268</u> -270, 271-273, <u>384, 385</u> 445-447- <u>450-468</u> -472-474, <u>497-505-519, 551, 552</u>

Tseng et al. Adeno-Associated Virus Serotype 1 (AAV1)- and AAV5-Antibody Complex Structures Reveal Evolutionary Commonalities in Parvovirus Antigenic Reactivity. Journal of Virology (2015) 89:1794-1808.

In the publication cited below multiple contact sites and footprint residues have been identified for ADK1a, that are very likely to be part of the binding site. The amino acids of each binding site are located in different parts of the protein chains and are recognized as the epitope of the antibody only in the assembled capsid where they are in close proximity to each other and in the correct conformation.

\*The residues in the VR are underlined. Those involved in AAV1 transduction are bold and italicized.

Tseng et al. Adeno-Associated Virus Serotype 1 (AAV1)- and AAV5-Antibody Complex Structures Reveal Evolutionary Commonalities in Parvovirus Antigenic Reactivity. Journal of Virology (2015) 89:1794-1808.



Neutralization of AAV6 with mouse monoclonal AAV1 antibody clone ADK1a (A) and human chimeric AAV1 antibody clone ADK1a-h1 (B) by using AAV6-NanoLuc<sup>®</sup> viral particles from Promega. (A) anti-AAV1 (intact particle) mouse monoclonal, ADK1a (Cat. No. 610150) or (B) anti-AAV1, human chimeric, ADK1a-h1 (Cat. No. 692350) were preincubated with AAV6-NanoLuc<sup>®</sup> viral particles for 30 min at RT at 300 rpm (antibody PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

concentrations 0.2-3,000 ng/ml). HEK293 cells (100 µl) were plated at 200,000 cells/ml in DMEM + 1% FCS. Virus-antibody-mix (20 µl) was added to the cells and incubated for 16-24 h at 37°C. Extracellular NanoLuc Inhibitor and Nano-Glo® Live Cell Assay System (Promega) was added to the wells and incubated for 5 min at RT at 300 rpm. Luminescence was measured using an ID5-Reader and plotted with Softmax Pro 7.1 software to determine the EC50 values.

## References

Publication	Species	Application
<a href="#">Sonntag, F. et al. The Assembly-Activating Protein Promotes Capsid Assembly of Different Adeno-Associated Virus Serotypes. J. Virol. 85, 12686â€“12697 (2011).</a>	AAV1	ELISA,ICC-IF
<a href="#">Adachi, K., Enoki, T., Kawano, Y., Veraz, M. &amp; Nakai, H. Drawing a high-resolution functional map of adeno-associated virus capsid by massively parallel sequencing. Nat. Commun. 5, (2014).</a>	AAV1	Neutralization epitope mapping
<a href="#">Tseng, Y.-S. et al. Adeno-Associated Virus Serotype 1 (AAV1)-and AAV5-Antibody Complex Structures Reveal Evolutionary Commonalities in Parvovirus Antigenic Reactivity. J. Virol. 89, 1794â€“1808 (2015).</a>	AAV1	epitope mapping,neutralization
<a href="#">Ohba K. et al. Adeno-associated virus vector system controlling capsid expression improves viral quantity and quality., iScience, 26, 106487, (2023).</a>	AAV1	IP
<a href="#">Kuck, D., Kern, A. &amp; Kleinschmidt, J. A. Development of AAV serotype-specific ELISAs using novel monoclonal antibodies. J. Virol. Methods 140, 17â€“24 (2007).</a>	AAV1,AAV6	ELISA,dot blot,IP,ICC-IF