

## Product datasheet

### protag-HiSec anti-chicken IgY-X2 AF568

#### Short overview

<b>Cat. No.</b>	80411
<b>Quantity</b>	500 µl

#### Product description

<b>Host</b>	Llama/alpaca
<b>Antibody Type</b>	Recombinant, produced in E.coli
<b>Isotype</b>	Single-domain antibody
<b>Clone</b>	3G2
<b>Immunogen</b>	Chicken IgY
<b>Formulation</b>	5 µM fluorescently labeled single-domain antibody in buffered saline, 50% glycerol, 0.09% sodium azide
<b>Note</b>	Centrifuge prior to opening
<b>Conjugate</b>	AF568
<b>Purification</b>	Affinity chromatography
<b>Storage</b>	Up to 3 months: -20°C; up to 12 months: -80°C or below; protect from light!
<b>Intended use</b>	Research use only
<b>Application</b>	ICC/IF, IHC
<b>Reactivity</b>	Chicken IgY
<b>No reactivity</b>	Goat IgG, Guinea pig IgG, Mouse IgG, Rabbit IgG, Rat IgG


#### Applications

<b>Immunocytochemistry (ICC)</b>	1:500
<b>Immunohistochemistry (IHC) - frozen</b>	1:500
<b>Immunohistochemistry (IHC) - paraffin</b>	1:500


#### Background

protag-HiSec anti-chicken IgY is an isotype- and species-specific camelid single-domain antibody (sdAb) produced by NanoTag Biotechnologies GmbH. It is directed against chicken IgY and contains two site-specifically coupled fluorophores per protag molecule. One single-domain antibody targets two fluorophores to a primary chicken antibody. Due to the monovalent binding, there are no primary and secondary antibody clusters formed, leading to better epitope accessibility and a more precise localization. There are no batch-to-batch variations, since the protag-HiSecs single-domain antibodies are recombinantly expressed.


#### Product images



Indirect immunostaining of PFA fixed mouse brain section with chicken anti-MAP2 and protag-HiSec anti-chicken IgY-X2 Sulfo-Cyanin 5 (green). Nuclei have been visualized by DAPI staining (blue)(courtesy of NanoTag Biotechnologies GmbH).



Mouse brain section with chicken anti-MAP 2 and protag-HiSec anti-chicken IgY-X2 Sulfo-Cyanin 5 (Cy5)(green). Nuclei were stained using DAPI (blue)(courtesy of NanoTag Biotechnologies GmbH).



Mouse cerebellum section stained with chicken anti-MAP 2 and protag-HiSec anti-chicken IgY-X2 Sulfo-Cyanin 5 (Cy5)(green). Nuclei were stained using DAPI (blue)(courtesy of NanoTag Biotechnologies GmbH).