

## Product datasheet

### anti-Plasma Cell Marker mouse monoclonal, EBS-O-231, purified

#### Short overview

|                      |                  |
|----------------------|------------------|
| <b>Cat. No.</b>      | 691670           |
| <b>Quantity</b>      | 1 ml (100 µg/ml) |
| <b>Concentration</b> | 100 µg/ml        |

#### Product description

|                      |                                       |
|----------------------|---------------------------------------|
| <b>Host</b>          | Mouse                                 |
| <b>Antibody Type</b> | Monoclonal                            |
| <b>Isotype</b>       | IgG2a kappa                           |
| <b>Clone</b>         | EBS-O-231                             |
| <b>Immunogen</b>     | Pancreatic cancer related serum mucin |
| <b>Formulation</b>   | PBS with 0.02% sodium azide           |
| <b>Conjugate</b>     | Unconjugated                          |
| <b>Purification</b>  | Affinity chromatography               |
| <b>Storage</b>       | 2-8°C                                 |
| <b>Intended use</b>  | Research use only                     |
| <b>Application</b>   | IHC                                   |
| <b>Reactivity</b>    | Human                                 |
| <b>No reactivity</b> | Rat                                   |

#### Applications

|  |  |
|--|--|
| <b>Immunohistochemistry (IHC) - paraffin</b> | 1:50-1:100 (1-2 µg/ml; microwave treatment in 10 mM citrate buffer pH 6.0 recommended) |
|--|--|

#### Background

EBS-O-231 recognizes an intra-cytoplasmic antigen, which shows a very high degree of specificity for plasma cells. This antigen is present in normal as well as neoplastic plasma cells. Plasma cells, which are large lymphocytes derived from an antigen-specific B cell, secrete antibodies and are responsible for humoral immunity. Plasma cells differentiate from B cells upon stimulation by CD4+ lymphocytes. The B cell acts as an antigen-presenting cell (APC), consuming an offending pathogen, which is taken up by the B cell by phagocytosis and broken down within proteosomes. Plasma cells contain basophilic cytoplasm; their nucleus contains heterochromatin organized in a characteristic cartwheel arrangement. This MAb superbly recognizes normal and neoplastic plasma cells in routine formalin-fixed, paraffin-embedded tissue sections. It is of potential value in identifying myeloma or plasmacytoma in bone marrow or other tissues. It also helps differentiate lympho-plasmacytoid lymphoma from lymphocytic and follicular lymphoma.


Positive control: human tonsil.

#### Product images

PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: [info@progen.com](mailto:info@progen.com) | Web: [www.progen.com](http://www.progen.com)

2024 April 26 / Version: 691670/DS-230420ibg | Page 1



Human tonsil