

Product datasheet

anti-Kappa Light Chain mouse monoclonal, EBS-huKappa, purified

Short overview

Cat. No.	691502
Quantity	1 ml (100 µg/ml)
Concentration	100 µg/ml

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1 kappa
Clone	EBS-huKappa
Immunogen	B lymphoma cells
Formulation	PBS with 0.02% sodium azide
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	2-8°C
Intended use	Research use only
Application	ELISA, FACS, ICC/IF, IHC
Reactivity	Human
No reactivity	Rat

Applications

ELISA	1:1-1:100,000 (0.001-100 µg/ml)
Flow Cytometry (FACS)	0.5-1.0 µg/million cells in 0.1 ml
Immunocytochemistry (ICC)	1:100-1:200 (0.5-1.0 µg/ml)
Immunohistochemistry (IHC) - frozen	1:50-1:100 (1-2 µg/ml)
Immunohistochemistry (IHC) - paraffin	1:50-1:100 (1-2 µg/ml; microwave treatment recommended)

Background

EBS-huKappa reacts with kappa light chain in mammals, the two light chains in an antibody are always identical, with only one type of light chain, kappa or lambda. The ratio of kappa to lambda is 70:30. However, with the occurrence of multiple myeloma or other B-cell malignancies this ratio is disturbed. Antibody to the kappa light chain is reportedly useful in the identification of leukemias, plasmacytomas, and certain non-Hodgkin's lymphomas. Demonstration of clonality in lymphoid infiltrates indicate that the infiltrate is malignant.

Positive control: 293T, Raji or hPBL cells, tonsil or spleen.

Product images



anti-Kappa Light Chain mouse monoclonal, EBS-huKappa, purified