

Product datasheet

AAV8 standard material (eGFP)

Short overview

Cat. No.	66V081
Quantity	100 µl
Concentration	> 1.0E+12 vg/ml; please find the lot-specific concentration (total capsid and viral genome titer) on the CoA

Product description

Filling grade	> 70% full capsids, please find the lot-specific filling grade on the CoA
Formulation	PBS + 0.014% Tween20 + 1 mM MgCl ₂ + 2.5 mM KCl
Source	Produced in HEK293T cells
Purity	> 95% by SDS-PAGE using Stain-Free Technology (Bio-Rad)
Quality check	Total capsid titer was assigned using AAV8 ELISA kits (PRAAV8 and PRAAV8XP, calibrated with ATCC reference standard material) and viral genome titer by qPCR using eGFP primers; QC included analysis of filling grade, purity, aggregation, and endotoxin testing.
Transfer Vector	pAAV-CMV-eGFP-WPRE
Packaging Plasmid	pRep2-Cap8 + pHelper
Endotoxin	< 1.0 EU/ml (detection limit 1.0 EU/ml)
Note	Please centrifuge before opening to ensure complete recovery of vial contents; aliquoting and repeated freeze/thaw cycles can lead to a drop in titer
Purification	Affinity Chromatography (POROS CaptureSelect AAVX Affinity Resin, Thermo Fisher Scientific), Iodixanol gradient centrifugation for removal of empty capsids using OptiPrep (Cat. No. 1893, PROGEN)
Storage	Up to 2 weeks: 2-8°C; long term storage in aliquots at -80°C; avoid > 5 freeze/thaw cycles
Intended use	Research use only
Application	Calibration of instruments e.g. mass photometry, Cell-based assay, Dot blot, ELISA, WB, ddPCR, qPCR

Applications

Dot Blot	Depending on primary antibody and detection method
ELISA	As a positive control in ELISA, dilute in ASSB 1x (provided with PROGENs AAV8 ELISA), analysis at least in duplicates is recommended
Cell-based Assay	Depending on the experimental setup
PCR	As standard or positive control in qPCR or ddPCR - concentration depending on experimental setup
Western Blot (WB)	Depending on primary antibody and detection method

Background

Our AAV8 standard material consists of fully assembled AAV8 capsids with an eGFP reporter gene controlled by a CMV promotor. Final

PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: info@progen.com | Web: www.progen.com


2025 February 15 / Version: 66V081/DS-171024ibg | Page 1

concentrations (total capsids and viral genome titer) are lot-specific and can be found on the CoA.


The AAV8 standard material is provided with a titer above $1.0E+12$ viral genomes/ml in a liquid formulation. Since the buffer does not contain any stabilizing proteins or dyes, the capsids can be used in various applications, including dot blot, western blot, ELISA, qPCR, ddPCR and cell-based assays. Our material contains only a very low remaining amount of iodixanol allowing DLS-SLS-UV/Vis analysis. The lot-specific titers were assigned using AAV8 ELISA (PRAAV8 and PRAAV8XP) and PCR using eGFP primers. Internal standard material calibrated with the ATCC reference standard material for AAV8 (VR-1816) was used for the final titer declaration. Our comprehensive quality control ensures well-characterized capsid material, which can be implemented as reference material in a variety of assays to prove the validity of the corresponding assay.

PROGEN provides eGFP-filled AAV standard material for the AAV serotypes 1, 2, 3, 5, 6, 8, 9, rh10, and rh74.

Product images



Transduction assay of AAV8 standard material (eGFP) using HeLa cells. HeLa cells ($50 \mu\text{l}$) were plated in 96-well plates at 200,000 cells/ml in DMEM + 10% FCS. $50 \mu\text{l}$ of eGFP-filled AAV8 capsids ($3.4E+10$ vg/ml) were added to the cells and incubated for 48 h at 37°C and 5% CO_2 . After incubation GFP-expression was visualized with a fluorescence microscope.



SDS PAGE with AAV8 eGFP-filled capsids. The AAV8 VP1, VP2 and VP3 proteins were separated on a 10% SDS PAGE and visualized by Pierce Silver stain kit. Only VP1, VP2 and VP3 proteins in the correct stoichiometry of 1:1:10 are detectable indicating a purity of the AAV preparation of >95%.



AAV8 standard material (eGFP)