

## Product datasheet

### anti-Myeloid Specific Antigen mouse monoclonal, BM-2, purified

#### Short overview

<b>Cat. No.</b>	691535
<b>Quantity</b>	1 ml (100 µg/ml)
<b>Concentration</b>	100 µg/ml

#### Product description

<b>Host</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Clone</b>	BM-2
<b>Immunogen</b>	Nuclei from pokeweed mitogen stimulated human peripheral blood lymphocytes
<b>Formulation</b>	PBS with 0.02% sodium azide
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity chromatography
<b>Storage</b>	2-8°C
<b>Intended use</b>	Research use only
<b>Application</b>	FACS, IHC
<b>Reactivity</b>	Human, Monkey

#### Applications


<b>Flow Cytometry (FACS)</b>	0.5-1.0 µg/million cells in 0.1 ml
<b>Immunohistochemistry (IHC) - frozen</b>	1:50-1:100 (1-2 µg/ml)
<b>Immunohistochemistry (IHC) - paraffin</b>	1:50-1:100 (1-2 µg/ml; microwave treatment in 10 mM citrate buffer pH 6.0 recommended)

#### Background

Until recently, immunological markers for myeloid cells have been lacking, especially those which identify different levels of cellular differentiation. The BM series provides a new panel of monoclonal antibodies which stain early precursor and mature forms of human myeloid cells. This panel of monoclonal antibodies reacts with antigenic determinants present in normal myeloid cells and leukemias of similar derivation. BM-2 recognizes a cytoplasmic antigen expressed in mature human granulocytes (polys) residing in lymphoid and non-lymphoid tissues. It does not react with any other cell type in human tissues.

Positive control: tonsil.

#### Product images



Human tonsil