

Product datasheet

anti-CD147 mouse monoclonal, CB-43, purified

Short overview

Cat. No.	691615
Quantity	1 ml (100 µg/ml)
Concentration	100 µg/ml

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgM kappa
Clone	CB-43
Immunogen	Human native leucocytes
Formulation	PBS with 0.02% sodium azide
UniprotID	P35613 (Human)
Synomym	Basigin, 5F7, Collagenase stimulatory factor, Extracellular matrix metalloproteinase inducer,
	EMMPRIN, Hepatoma-associated antigen, HAb18G, Leukocyte activation antigen M6, OK blood
	group antigen, Tumor cell-derived collagenase stimulatory factor, TCSF, CD antigen CD147,
	BSG, UNQ6505/PRO21383
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	2-8°C
Intended use	Research use only
Application	FACS, ICC/IF, IHC
Reactivity	Human

Applications

Flow Cytometry (FACS) Immunocytochemistry (ICC) Immunohistochemistry (IHC) - frozen

0.5-1.0 µg/million cells in 0.1 ml 1:100-1:200 (0.5-1.0 µg/ml) 1:50-1:100 (1-2 µg/ml)


Background

CD147 is a transmembrane glycoprotein of the immunoglobulin superfamily. It is expressed more intensely on thymocytes than on mature peripheral blood T cells. CD147 is important in spermatogenesis, embryo implantation, neural network formation, and tumor progression. CD147 is involved in the regulation of matrix remodeling at the epidermal-dermal interface. It stimulates the production of interstitial collagenase, gelatinase A, stromelysin-1 and various metalloproteinases (MMPs) by fibroblasts. These enzymes, which are typically increased during tissue degradation and wound healing, are important factors in cancer invasion and metastasis.

Positive control: HSB2 cells, renal cell, ovarian carcinoma or melanoma.

PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: info@progen.com | Web: www.progen.com 2024 April 23 / Version: 691615/DS-230420ibg | Page 1

Product images



Human kidney